

## ISOLATION AND MOLECULAR IDENTIFICATION OF FUNGAL ISOLATES FROM STORED CEREALS USING PCR-RFLP METHOD

Roxana Zaharia<sup>1</sup>, Cristina Petrișor<sup>1\*</sup>, Petruța Cornea<sup>2</sup>, Camelia Diguță<sup>2</sup>,  
Stelica Cristea<sup>2</sup>, Ștefan Sorin<sup>1</sup>

<sup>1</sup>Research and Development Institute for Plant Protection Bucharest, 8 Ion Ionescu de la Brad Blvd,  
District 1, Bucharest, Romania

<sup>2</sup>University of Agronomic Sciences and Veterinary Medicine of Bucharest, 59 Mărăști Blvd,  
District 1, Bucharest, Romania

\*Corresponding author. E-mail: crisstop@yahoo.com

### ABSTRACT

Contamination of grain cereals with toxic metabolites of fungi, both pathogenic and saprotrophic, is one of the particularly important problems in global agriculture. The aim of the current study was molecular identification of fungi isolates from different samples of stored cereals and evaluate the utility of PCR-RFLP of the ITS region technique. The results established that the most abundant species were found belonging to *Aspergillus* genera (50%), followed by *Fusarium* spp. (19%) and *Penicillium* spp. (19%). *Aspergillus flavus* was the most frequent species, representing almost 40% of the isolates belonging to the genus *Aspergillus*. Also were identified as *Aspergillus versicolor*, *Aspergillus ruber* and *Aspergillus niger* by molecular analysis representing 10% each.

**Keywords:** wheat, electrophoresis, fungal contamination, DNA amplification, DNA sequencing.

### INTRODUCTION

Cereals are natural substrates for fungal development due to their nutritional composition rich in starch and protein, proving necessary carbohydrate and nitrogen for development of micromycetes thus contamination leading to financial and food safety losses (Fleurat-Lessard, 2017).

Food grains may be stored at different levels in the food supply chain in silos, warehouses, bags, containers, traditional storage structures, or in other defined units. The occurrence of molds in stored grains is influenced by a range of abiotic and biotic factors like temperature, water availability and intergranular gas composition, contaminant molds, insect pests, rodents and preservatives which are added to conserve moist grain for animal feed (Magan et al., 2003; Fleurat-Lessard, 2017) and represents an important problem in different countries worldwide.

Many species of fungi that are found on the surface or inside grains at harvest may cause spoilage in stored grains and produce

mycotoxins that severely decrease crop value and are harmful to animal and human health that consume the contaminated feed or grain products (Placinta et al., 1999; Magan et al., 2010; Shiju et al., 2010; Gonzales Perreyra et al., 2011; Belkacem-Hanfi et al., 2013; Jedidi et al., 2018; Kumari et al., 2019).

Studies on the several filamentous fungi (*Aspergillus*, *Fusarium*, *Penicillium*, *Alternaria*) which attacked stored cereal grain and are responsible for production of toxic metabolites for humans and animals (Belkacem-Hanfi et al., 2013; Comby et al., 2016; Gdanetz and Trail, 2017) are well documented.

These fungi can be identified by traditional plating techniques based on the morphological characteristics such as the shape and the size of the conidia, the presence/absence of chlamydospores and also the colony morphology. However conventional methods used for fungal detection are labour and time-consuming require mycological expertise and not always very specific (Jurado et al., 2006) In general, it is difficult to distinguish between species having similar

morphological characteristics, particularly when using traditional methods.

Thus the molecular approaches have been developed for toxigenic fungi studies like polymerase chain reaction (PCR) techniques which are rapid, sensitive, specific and reliable diagnosis methods in species identification without additional confirmation steps (Nicholson et al., 1998; Jurado et al., 2006; Suanthie et al., 2009; Datta et al., 2011; Atoui et al., 2011; Kumari et al., 2016; Sadhasivam et al., 2017; Jayachandran et al., 2019).

However, a widely used method is PCR-RFLP a powerful technique generally used for microbial identification and classification even at species level, by which PCR amplification of a target region is performed, followed by digestion of the fragment amplified with restriction enzymes. Separation of the obtained fragments is performed by electrophoresis to visualize the differences between the restrictions profiles obtained.

This technique was used for the rapid identification of the large number of fungi (Somashekar et al., 2004; Dupont et al., 2006; Llorens et al., 2006; González-Salgado et al., 2009; Diguță et al., 2011; Kizis et al., 2014; Ahmad et al., 2014; Kachuei et al., 2015). No studies have been reported in Romania on the use of the PCR-RFLP method for identification of fungitoxic species from stored grains.

The studies of the Romanian authors were focused on the analysis of the cereals mycoflora (Tabuc et al., 2009; Cornea et al., 2013; Misca et al., 2014; Dudoiu et al., 2016; Stanciu et al., 2017) and their content of mycotoxins (Placinta et al., 1999; Alexa et al., 2013; Misca et al., 2014; Bozac et al., 2016; Gagiú et al., 2018).

In light of these informations, the objective of present study was to examine the fungal contamination of different stored wheat samples using molecular tests species specific RFLP-PCR assays.

## MATERIAL AND METHODS

Two hundred samples of contaminated wheat were collected from different grain

warehouse located in different areas of Păulești, Prahova County in the years 2015-2017. From each warehouse a sample of 1 kg was collected from the front, from the center and from different points in the horizontal depth of the grain mass and also in the area close to the walls of the warehouse.

The water content of cereal seeds varied between 10-14%. The collected samples were kept in sterile plastic bags during transport and 100 g of each sample was frozen in liquid nitrogen, lyophilized and ground in a fine puberty. The samples were kept at 4°C until analysis.

### 1. Fungal DNA extraction

The fungal isolates were grown in PDB (Potato Dextrose Broth, VWR Chemicals) liquid medium, at a temperature between 28-30°C, for 96 hours. The fungal biomass was treated with liquid nitrogen in order to break down the cell wall. Subsequently, the fungal suspension obtained was centrifuged at 10.000 rpm for 10 minutes, thus resulting a sediment of approximately 100-150 mg. The obtained sediment was used as a base for fungal DNA extraction using the ZR Fungal/Bacterial MiniPrep kit (Zymo Research, USA).

Following the protocol, 750 µl of lysis solution was added to the fungal sediment and the obtained mixture was vortexed at maximum speed for 5 minutes. The obtained suspension was centrifuged at 10.000 x g for 1 minute; The binding buffer (Fungal/Bacterial DNA Binding Buffer) - supplemented with 0.05% beta-mercaptoethanol (v/v) was added to the filtered suspension; the resulting mixture was centrifuged at 10000xg for 1 minute. A pre-wash buffer (DNA Pre-Wash buffer) was added and a new centrifugation was performed. Next, wash buffer (Fungal/Bacterial DNA Wash Buffer) was added to the supernatant and again the mixture was centrifuged 10000xg for 1 minute. In the final step, the DNA was resuspended in 75 µl of DNA elution buffer.

### 2. DNA amplification and sequencing

RFLP-PCR-ITS specific assays were performed using ITS1 (5'-

TCCGTAGGTGAACCTGCGG-3') and ITS4 (5'-TCCTCCGCTTATTGATATGC-3') amplification primers, retained to amplify the 5.8S-ITS region.

PCR amplification reactions were performed in a final volume of 50  $\mu$ l containing 10X DreamTaq Green Buffer (containing 20 mM MgCl<sub>2</sub>), 100-150 ng/DNA, 10 mM dNTP, 10  $\mu$ M from each primer and DNA polymerase 5 U/ $\mu$ L. The reaction mixture was vortexed and then subjected to PCR reaction.

PCR amplification was performed in a thermal cycler (Multi Gene, Labnet), the

reaction mixture being subjected to several thermocycles with the following profile: 1 cycle representing the initial denaturation (3 minutes at 94°C), 35 cycles consisting of denaturation (1.5 minutes at 94°C), hybridization (1.5 minutes at 55.5°C) and elongation (2 minutes at 72°C), followed by 1 cycle for final elongation (10 minutes at 72°C) and then the final cycle at 40°C.

Regarding the digestion of PCR products of the fungal isolates obtained it was performed using the following restriction enzymes (Thermo Fisher Scientific):

| <i>Cfr9I</i> :  | <i>HaeIII</i> :  | <i>HinfI</i> :   | <i>MseI</i> :  | <i>SduI</i> :  |
|---|--|--|--|--|
| 5'...C <sup>^</sup> CCGGG...3'<br>3'...GGCC <sup>^</sup> C...5' | 5'..GG <sup>^</sup> CC..3'<br>3'..CC <sup>^</sup> GG..5' | 5'...G <sup>^</sup> ANTC...3'<br>3'...CTNA <sup>^</sup> G...5' | 5'...T <sup>^</sup> TAA...3'<br>3'...AAT <sup>^</sup> T...5' | 5'...GdGCh <sup>^</sup> C...3'<br>3'...C <sup>^</sup> hCGdG...5' |

The reaction mixture consists of PCR product, phosphate buffer, nuclease-free water and restriction enzymes. The enzymatic digestion was performed on a water bath during 4 hours at 37°C for SduI, HinfI; HaeIII; Cfr9I and at 65°C for the enzyme MseI.

### 3. DNA electrophoresis

The quality of the isolated fungal DNA was verified using agarose gel electrophoresis. The samples were loaded into a 2% agarose gel prepared in 1X TBE buffer (Tris-Borate-EDTA) containing 10 mg/mL ethidium bromide. 8  $\mu$ l of PCR product, respectively 10 -15  $\mu$ l of restriction product were charged and the electrophoresis was performed using 1X TBE buffer at 90 V, for 1 hour until the front of the blue bromophenol dye migrated near the end of the gel. The molecular standard used was DNA Ladder 100 bp (GeneRuler 100bp Plus DNA Ladder, Thermo Scientific, USA).

Subsequently, PCR and restriction products were visualized using a UV transilluminator

(GelDoc-IT Imaging Systems) by exposing ethidium bromide to UV light ( $\lambda$ =250-310 nm). The dimensions of the PCR products and the restriction fragments obtained were estimated by comparison with a standard length of DNA (GeneRuler 100bp DNA Ladder, Thermo Fisher Scientific). The analysis of the obtained restriction profiles was performed by comparing with the restriction profiles (theoretically obtained using the program <http://biotools.umassmed.edu/tacg4/>) the sequences already existing in the National Center for Biotechnology Information database (<https://www.ncbi.nlm.nih.gov/nucleotide/>).

## RESULTS AND DISCUSSION

The initially performed morphological analysis (macroscopic and microscopic) on the 16 studied fungal isolates showed that the mycotoxigenic microflora detected on stored cereals is dominated by three important genera: *Aspergillus*, *Fusarium*, *Penicillium* (Figure 1).




Figure 1. Isolated fungal strains on PDA media

a. *A. niger*; b. *A. parasiticus*; c. *A. flavus*; d. *Fusarium* sp.; e. *Penicillium* sp.

However, classical methods are not 100% efficient and at the same time they require high experience of researchers in terms of mold taxonomy. Thus, it is important that these mycotoxic molds to be identified quickly and accurately to ensure the safety of consumers and animals. Given that, an alternative to conventional microbiological techniques for diagnosing phytopathogenic

fungi is represented by PCR molecular methods.

Efficient identification and discrimination of fungal species was performed with the PCR-RFLP method by selecting the 5,8S-ITS region. Following the amplification using the universal primers ITS1/ITS4 there were obtained PCR products with dimensions of approximately 540-600 bp (Figure 2).




Figure 2. Electrophoresis in 2% agarose gel of ITS1/ITS4 amplified products for the fungal isolates **1 - MR2; 2 - Tr; 3 - P3; 4 - Pg; 5 - D1; 6 - Av; 7 - Px; 8 - P1; 9 - P2; 10 - Piz; 11 - Rx; 12 - SS; 13 - An; 14 - F1; 15 - F2; 16 - F3; M - negative control; L - GeneRuler 100 bp Plus DNA Ladder.**

Some authors claim that a single amplification by PCR-ITS only allows the identification/differentiation of fungal isolates to a small extent at the species level (Diguță et al., 2011). The obtained PCR products were subjected to separate digestion with the

restriction enzymes SduI and HinfI. A third MseI or HaeIII enzyme was used to complete species-level identification. The resulting fragments from the enzymatic cleavage were visualized in agarose gel electrophoresis and are shown in Figures 3 and 4.

ROXANA ZAHARIA ET AL.: ISOLATION AND MOLECULAR IDENTIFICATION OF FUNGAL ISOLATES FROM STORED CEREALS USING PCR-RFLP METHOD




Figure 3. Restriction profile of the studied fungal isolates, after restriction endonucleases digestion:

A) *Sma*I; B) *Hinf*I; C) *Mse*I

1 - MR2; 2 - Tr; 3 - P3; 4 - Pg; 5 - DI; 6 - Av; 7 - Px; 8 - PI; 9 - P2; 10 - Piz; 11 - Rx; 12 - SS; 13 - An;  
L - GeneRuler 100 bp Plus DNA Ladder




Figure 4. *Fusarium* spp. restriction profile after digestion with restriction endonucleases *SduI*; *HinfI*; *HaeIII* 14 - F1; 15 - F2; 16 - F3; L - GeneRuler 100 bp Plus DNA Ladder.

The restriction profiles obtained with each restriction enzyme are summarized in Table 1.

Table 1. Identification with PCR-ITS-RFLP of studied isolated fungi

| Nr. crt. | Fungal isolate | Restriction fragments (bp) |              |             |               | Identification                 |
|----------|----------------|----------------------------|--------------|-------------|---------------|--------------------------------|
|          |                | <i>SduI</i>                | <i>HinfI</i> | <i>MseI</i> | <i>HaeIII</i> |                                |
| 1.       | MR2            | 110 170 300                | 110 170 300  | 560         |               | <i>Aspergillus versicolor</i>  |
| 2.       | Tr             | 190 400                    | 280 300      | 500         |               | <i>Trichoderma viride</i>      |
| 3.       | P3             | 170 300                    | 300 300      | 210 370     |               | <i>Aspergillus flavus</i>      |
| 4.       | Pg             | 170 300                    | 300 300      | 210 370     |               | <i>Aspergillus flavus</i>      |
| 5.       | D1             | 170 300                    | 300 300      | 210 370     |               | <i>Aspergillus flavus</i>      |
| 6.       | Av             | 170 300                    | 300 300      | 210 370     |               | <i>Aspergillus flavus</i>      |
| 7.       | Px             | 110 170 300                | 110 170 300  | 560         |               | <i>Aspergillus versicolor</i>  |
| 8.       | P1             | 170 270                    | 110 190 300  | 110 370     |               | <i>Penicillium expansum</i>    |
| 9.       | P2             | 170 270                    | 300 300      | 110 370     |               | <i>Penicillium chrysogenum</i> |
| 10.      | Piz            | 170 270                    | 110 190 300  | 110 370     |               | <i>Penicillium expansum</i>    |
| 11.      | Rx             | 170 340                    | 280 280      | 180 280     |               | <i>Aspergillus ruber</i>       |
| 12.      | SS             | 540                        | 180 280      | 110 430     |               | <i>Botrytis cinerea</i>        |
| 13.      | An             | 170 300                    | 300 300      | 210 300     |               | <i>Aspergillus niger</i>       |
| 14.      | F1             | 560                        | 280 280      |             | 100 120 360   | <i>Fusarium tricinctum</i>     |
| 15.      | F2             | 180 380                    | 280 300      |             | 80 110 280    | <i>Fusarium proliferatum</i>   |
| 16.      | F3             | 560                        | 120 170 280  |             | 560           | <i>Fusarium</i> sp.            |

The resulting restriction profiles were compared with the restriction profiles of the sequences available in the theoretically obtained NCBI database.

Among *Aspergillus* sp. identified in this study, four isolates showed a specific profile of *A. flavus*, two isolates a specific profile of *A. versicolor*. One isolate was identified as *A. niger* and one isolate as *A. ruber*. Similar

results regarding the dominance of the genus *Aspergillus* in the stored wheat samples were also presented in the studies of some Moroccan and Indian authors (Hajjaji et al., 2006; Kumari and Ghosh, 2016). Also the results of Jedidi et al. (2018) established that the wheat samples were contaminated especially with *Aspergillus* but also with *Alternaria* and *Eurotium*. However,

Algerian researchers have also identified *A. orchraceus* (Riba et al., 2008).

Ahmad et al. (2014) differentiated *A. flavus*, known as a aflatoxin producer, from *A. parasiticus* based on the restriction profiles obtained with the BanI and NlaIV enzymes of the amplified product from the *aflP* gene. Also, Somashekar et al. (2004) differentiated *A. flavus* from *A. parasiticus* based on restriction profiles of aflR amplicons with the enzyme PvuII. Specific detection of aflatoxin molds was performed on wheat samples using specific primers for the internally transcribed region (ITS1-5.8S-ITS2) (González-Salgado et al., 2009; Sardinãs et al., 2010).

The second major fungal genus was *Penicillium*. For the differentiation of *Penicillium* isolates, the restriction profiles obtained were compared with those obtained by Diguță et al. (2011). According to the results presented in Table 1, two isolates were identified as *P. expansum* and one isolate had a specific profile for *P. chrysogenum*/*P. crustosum*/*P. commune*. In this case, the Cfr9I endonuclease was used to complete the identification, and the resulting restriction profile corresponds to the *P. chrysogenum* species. Dupont et al. (2006) developed a PCR-RFLP method using a combination of 5 endonucleases that allowed the differentiation of 12 species of *Penicillium* from the *Biverticillium* subgenus. In another study, Diguță et al. (2011) optimized a PCR-ITS-RFLP method for the identification and differentiation of 22 different species out of a total of 24 *Penicillium* species detected on grape berries, with the exception of *P. thomii* and *P. glabrum* which remained non-differentiable. Also, Rousseaux and Guilloux-Bénatier (2016) using the PCR-RFLP technique were able to differentiate 22 species of the genus *Penicillium*. Previous studies (Kumari and Ghosh, 2016) also reported the existence of fungal species of the genera *Aspergillus* and *Penicillium* in stored wheat samples.

The restriction profiles analysis for the 3 isolates belonging to the genus *Fusarium*, obtained after enzymatic digestion with SduI,

HinFI and HaeIII, allowed the identification of *F. tricinctum* and *F. proliferatum* species (based on the sequences already existing in the National Center for Biotechnology Information database (<https://www.ncbi.nlm.nih.gov/nucleotide/>)). For the final identification of the F3 isolate, an analysis of the restriction profiles of other *Fusarium* species will be made, and the validation of the results will be done by sequencing. Llorens et al. (2006) amplified the IGS gene followed by digestion with 6 restriction enzymes (CfoI, AluI, HapII, XhoI, EcoRI and PstI) to differentiate 6 species of *Fusarium* (*F. culmorum*, *F. graminearum*, *F. cerealis*, *F. poae*, *F. oxysporum*, and *Gibberella fujikuroi*). The results are consistent with those of the authors Sudharsan et al. (2017). Studies by Kachuei et al. (2015) showed that by using the restriction enzymes HhaI, MspI, several species of *Fusarium*, such as *F. dimerum*, *F. subglabratum*, *F. beomiform*, *F. equiseti* and *F. compactum* were differentiated. They also used MspI and TaqI enzymes to distinguish the following *Fusarium* species: *F. sacchari*, *F. brevicatenuatum*, *F. concolor* and *F. ambrosium*.

In this study, the most fungal strains identified by PCR-ITS-RFLP belong to the genus *Aspergillus* (50% of total fungal isolates), followed by *Fusarium* (19%) and *Penicillium* (19%). Also, were identified two isolates belonging to the species *Botrytis cinerea* and *Trichoderma* sp., respectively. The detection of these fungal species is of major importance, as most isolated species are potentially producing mycotoxins such as: aflatoxins, fumonisins, ochratoxin A, zearalenone and deoxynivalenol which represent a serious problem worldwide (FAO 2004), causing many negative effects on human and animal health.

The detection of different species belonging to the genera *Aspergillus* and *Penicillium* can be explained by improper storage of cereals (high humidity, fluctuating temperatures, lack of aeration), which favors the growth of these molds known as storage molds. *Aspergillus* sp. has been shown to be the most common causative agent of grain

rot, with *A. flavus* being the most predominant species detected. *Penicillium expansum* has been frequently detected on fruits (apples, grapes) (Diguță et al., 2011) and only occasionally in cereals.

*Fusarium* is one of the major toxin-producing genera, such as fumonisins and trichothecenes. *Fusarium* sp. are molds that frequently contaminate grains worldwide (Sampietro et al., 2011; Jedidi et al., 2018). *Fusarium* sp. are considered plant pathogens and are responsible for infections before and after harvest, however, some species of *Fusarium* are able to persist in harvested and stored grain and grow in warehouses when humidity conditions become favorable. *F. graminearum*, considered the main contaminant of wheat grains, was not detected in this study. At the same time, the studies of Sadhasivam et al. (2017) claim that the application of the multiplex PCR method on wheat samples stored in warehouses led to the detection of *Aspergillus* and *Fusarium* genera especially.

## CONCLUSIONS

RFLP-PCR methods have been successfully applied for the isolation and identification of pathogenic fungi that contaminate stored wheat, having the advantage of being fast and specific.

The fungal strains identified by PCR-ITS-RFLP belong to the genus *Aspergillus* (50% of total fungal isolates), followed by *Fusarium* (19%) and *Penicillium* (19%).

## REFERENCES

- Ahmad, M.M., Ahmad, M., Ali, A., Hamid, R., Javed, S., Abidin, M.Z., 2014. *Detection of Aspergillus flavus and Aspergillus parasiticus from aflatoxin-contaminated peanuts and their differentiation using PCR-RFLP*. Ann. of Microbiology, 64: 1597-1605.
- Alexa, E., Dehelean, C.A., Poiana, M.A., Radulov, I., Cimpean, A.M., Bordean, D.M., Tulcan, C., Pop, G., 2013. *The occurrence of mycotoxins in wheat from western Romania and histopathological impact as the effect of feed intake*. Chem. Cent. J., 7: 2-11.
- Atoui, A., El Khoury, A., Kallassy, M., Lebrihi, A., 2011. *Quantification of Fusarium graminearum and Fusarium culmorum by real-time PCR system and zearalenone assessment in maize*. Int. J. of Food Microbiol., 154(1-2): 59-65. ISSN 0168-1605
- Belkacem-Hanfi, N., Semmar, N., Perraud-Gaime, I., Guesmi, A., Cherni, M., Cherif, I., Boudabous, A., Roussos, S., 2013. *Spatio-temporal analysis of post-harvest moulds genera distribution on stored durum wheat cultivated in Tunisia*. Journal of Stored Product Research, 55: 116-123.
- Bozac, P., Popescu, S., Botau, D., Boldura, O.M., Pirvulescu, P., 2016. *Molecular characterization for some new Fusarium isolates collected from the West part of Romania*. Romanian Biotechnological Letters, 21(3): 11560-11568.
- Comby, M., Lacoste, S., Baillieul, F., Profizi, C., Dupont, J., 2016. *Spatial and temporal variation of cultivable communities of co-occurring endophytes and pathogens in wheat*. Front. Microbiol., 7: 403.
- Cornea, C.P., Isael-Roming, F., Ciucă, M., Voaideş, C., 2013. *Natural occurrence of Fusarium species and corresponding chemotypes in wheat scab complex from Romania*. Romanian Biotechnological Letters, 18(6): 8787-8795.
- Datta, S., Choudhary, R.G., Shamim, M.D., Vishwa, D., 2011. *Polymorphism in the internal transcribed spacer (ITS) region of the ribosomal DNA among different Fusarium species*. Archives of Phytopathology and Plant Protection, 44(6): 558-566
- Diguță, C.F., Vincent, B., Guilloux-Benatier, M., Alexandre, H., Rousseaux, S., 2011. *PCR ITS-RFLP: a useful method for identifying filamentous fungi isolates on grapes*. Food Microbiol., 28(6): 1145-1154.
- Dudoiu, R., Cristea, S., Lupu, C., Popa, D., Oprea, M., 2016. *Micoflora associated with maize grains during storage period*. Agrolife Scientific Journal, 5(1): 58-63.
- Dupont, J., Denetiere, B., Jacquet, C., Roquebert, M.F., 2006. *PCR-RFLP of ITS rDNA for the rapid identification of Penicillium subgenus Biverticillium species*. Revista Iberoamericana Micologia, 26: 145-150.
- Fleurat-Lessard, F., 2017. *Integrated management of the risks of stored grain spoilage by seedborne fungi and contamination by storage mould mycotoxins - An update*. Journal of Stored Products Research, 71: 22-40.
- Gagiu, V., Mateescu, E., Armeanu, I., Dobre, A.A., Smeu, I., Cucu, M.E., Oprea, O.A., Iorga, E., Belc, A., 2018. *Post-harvest contamination with mycotoxins in the context of the geographic and agroclimatic conditions in Romania*. Toxins, 10(12): 533.
- Gdanetz, K., and Trail, F., 2017. *The wheat microbiome under four management strategies, and potential for endophytes in disease protection*. Phytobiomes, 1: 158-168.
- Gonzalez Pereyra, M.L., Chiacchiera, S.M., Rosa, C.A.R., Sager Dalceron, A.M., Cavaglieri, L., 2011. *Comparative analysis of the mycobiota and*



ROXANA ZAHARIA ET AL.: ISOLATION AND MOLECULAR IDENTIFICATION OF FUNGAL ISOLATES FROM STORED CEREALS USING PCR-RFLP METHOD

- mycotoxins contaminating corn trench silos and silo bags*. J. Sci. Food. Agr., 91(8): 1474-1481.
- González-Salgado, A., Patiño, B., Gil-Serna, J., Vázquez, C., González-Jaén, M.T., 2009. *Specific detection of Aspergillus carbonarius by SYBR® Green and TaqMan® quantitative PCR assays based on the multicopy ITS2 region of the rRNA gene*. FEMS Microbiology Letters, 295: 57-66.
- Hajjaji, A., El Otmani, M., Bouya, D., Bouseta, A., Mathieu, F., Collin, S., Lebrihi, A., 2006. *Occurrence of mycotoxins (ochratoxin A, deoxynivalenol) and toxigenic fungi in Moroccan wheat grains: impact of ecological factors on the growth and ochratoxin A production*. Mol. Nutr. Food. Res., 50: 494-499.
- Jayachandran, L.E., Kumari, R., Rao, P.S., 2019. *Modulation of mycobiota associated with rice stored in bamboo reinforced concrete silo-molecular identification and phenotypic characterization*. Journal of Stored Products Research, 83: 14-24.
- Jedidi, I., Soldevilla, C., Lahouar, A., Marín, P., González-Jaén, M.T., Said, S., 2018. *Mycoflora isolation and molecular characterization of Aspergillus and Fusarium species in Tunisian cereals*. Saudi Journal of Biological Sciences, 25: 868-874.
- Jurado, M., Vázquez, C., Marín, S., Sanchi, V., González-Jaén, M.T., 2006. *PCR-based strategy to detect contamination with mycotoxigenic Fusarium species in maize*. Syst. Appl. Microbiol., 29: 681-689.
- Kachuei, R., Yadegari, M.H., Safaie, N., Ghiasian, A., Noorbakhsh, F., Piranfar, V., Rezaie, S., 2015. *PCR - RFLP patterns for the differentiation of the Fusarium species in virtue of ITS rDNA*. Curr. Med. Mycol., 1(1): 4-11.
- Kizis, D., Natskoulis, P., Nychas, G.J., Panagou, E.Z., 2014. *Biodiversity and ITS-RFLP characterisation of Aspergillus section Nigri isolates in grapes from four traditional grape-producing areas in Greece*. PLoS ONE, 9(4): 93923.
- Kumari, R., and Ghosh, A.K., 2016. *Molecular characterization of fungi present in stored food grains*. International Conference on Emerging Technologies in Agricultural and Food Engineering (ETAE), IIT Kharagpur. ISBN: 978-93-86256-30-0.
- Kumari, R., Jayachandran, L.E. Ghosh, A.K., 2019. *Investigation of diversity and dominance of fungal biota in stored wheat grains from governmental warehouses in West Bengal, India*. J. Sci. Food Agric., 99: 3490-3500.
- Llorens, A., Hinojo, M.J., Mateo, R., Gonzalez-Jaen, M.T., Valle-Algarra, F.M., Logrieco, A., Jimenez, M., 2006. *Characterization of Fusarium spp. isolates by PCR-RFLP analysis of the intergenic spacer region of the rRNA gene (rDNA)*. Int. J. Food Microbiol., 106: 297-306.
- Magan, N., Hope, R., Cairns, V., Aldred, D., 2003. *Post-harvest fungal ecology: impact of fungal growth and mycotoxin accumulation in stored grain*. European J. Plant Pathol., 109: 723-730.
- Magan, N., Aldred, D., Hope, R., Mitchell, D., 2010. *Grain and Grapes: effects on growth, deoxynivalenol and ochratoxin production by Fusarium culmorum and Aspergillus carbonarius*. Toxins, 2: 353-366.
- Misca, C. Daminescu, L., Jianu, C., David, I., Misca, L., Marginean, O., Radoi, B., Rinovetz, A, Bujanc, G., Velcirov, A., 2014. *The incidence of the strains of Fusarium sp. and of zearalenone in cereals analyzed from the south west of Romania*. Ann. West Univ. Timisoara Ser. Biol., XVII: 137-144.
- Nicholson, P., Simpson, D.R., Weston, G., Rezanoor, H.N., Lees, A.K., Parry, D.W., Joyce, D., 1998. *Detection and quantification of Fusarium culmorum and Fusarium graminearum in cereals using PCR assays*. Physiol. Mol. Plant Pathol., 53: 17-37.
- Placinta, C.M., D'Mello, C.P.F., MacDonald, A.M.C., 1999. *A review of worldwide contamination of cereal grains and animal feed with Fusarium mycotoxins*. Anim. Feed Sci. Technol., 78: 21-37.
- Riba, A., Mokrane, S., Mathieu, F., Lebrihi, A., Sabaou, N., 2008. *Mycoflora and ochratoxin. A producing strains of Aspergillus in Algerian wheat*. Int. J. Food Microbiol., 122: 85-92.
- Rousseaux, S., and Guilloux-Bénatier, M., 2016. *PCR ITS-RFLP for Penicillium species and other genera*. In: Moretti & Susca (eds.), Mycotoxigenic fungi - Methods and Protocols. Methods in Molecular Biology, 1542: 321-333.
- Sadhasivam, S., Britzi, M., Zakin, V., Kostyukovsky, M., Trostanetsky, A., Quinn, E., Sionov, E., 2017. *Rapid detection and identification of mycotoxigenic fungi and mycotoxins in stored wheat grain*. Toxins, 9(302): 1-17.
- Sampietro, D.A., Diaz, C.G., Gonzalez, V., Vattuone, M.A., Ploper, L.D., Catalan, C.A., Ward, T.J., 2011. *Species diversity and toxigenic potential of Fusarium graminearum complex isolates from maize fields in northwest Argentina*. Int. J. Food Microbiol., 145: 359-364.
- Sardiñas, N., Vázquez, C., Gil-Serna, J., González-Jaén, M.T., Patiño, B., 2010. *Specific detection of Aspergillus parasiticus in wheat flour using a highly sensitive PCR assay*. Food Add. Contamin., 27: 853-858.
- Shiju, M., Thomas, G., Ahmad, T., 2010. *An evaluation on the impact of fungi on the post-harvested stored wheat grains*. International Journal of Biotech. and Biochem., 6: 995-1002.
- Somashekar, D., Rati, E.R., Chandrashekar, A., 2004. *PCR-restriction fragment length analysis of aflR gene for differentiation and detection of Aspergillus flavus and Aspergillus parasiticus in maize*. Int. J. Food Microbiol., 93: 101-107.

- Stanciu, O., Juan, C., Miere, D., Loghin, F., Mañes, J., 2017. *Presence of enniatins and beauvericin in Romanian wheat samples: From raw material to products for direct human consumption*. *Toxins*, 9(189): 1-16.
- Suanthie, Y., Cousin, M.A., Woloshuk, C.P., 2009. *Multiplex real-time PCR for detection and quantification of mycotoxigenic Aspergillus, Penicillium and Fusarium*. *Journal of Stored Products Research*, 45: 139-145.
- Sudharsan, S., Malka, B., Varda, Z., Moshe, K., Anatoly, T., Elazar, Q., Edward, S., 2017. *Rapid detection and identification of mycotoxigenic fungi and mycotoxins in stored wheat grain*. *Toxins*, 9(10): 1-17.
- Tabuc, C., Marin, D., Guerre, P., Sesan, T., Bailly, J., 2009. *Mold and mycotoxin content of cereals in southeastern of Romania*. *Journal of Food Protection*, 72: 662-665.