

IMPACTS OF SILICON FOLIAR SPRAYING AND NITROGEN APPLICATION TECHNIQUES ON QUANTITATIVE AND QUALITATIVE PARAMETERS OF RICE AT DIFFERENT PLANTING SPACES

Sarvenaz Yahyazadeh, Hamidreza Mobasser*, Elyas Rahimi Petroudi, Alireza Daneshmand

Department of Agronomy, Qaemshahr Branch, Islamic Azad University, Qaemshahr, Iran

*Corresponding author. E-mail: drmobasser.neg@gmail.com

ABSTRACT

The silicon (Si) foliar application along with timely application of nitrogen (N) can be help to increase the quantity and quality of crops. Also, planting spaces is an agronomic technique that can affect yield and quality of rice. The objective of this study was to evaluate the effects of Si foliar application and N application techniques on yield components, grain yield and accumulation of Si and N in rice grains at different planting spaces. The experiment was conducted as a split-split-plot in a randomized complete block design with three replications during two crop years of 2018 and 2019. The experimental treatments included the main plot assigned to planting spaces (25×10 cm, 20×10 cm), the sub-plot assigned to N application techniques (N application as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading, N application as 40% at basal + 40% at panicle initiation + 20% at full heading), and the sub-sub-plot assigned to foliar application of Si (control or non-use of Si, Si foliar application). The results indicated that the reducing planting space (20×10 cm) resulted in an increase in the number of panicle number m⁻² by 9.3%, followed by an improvement in grain yield by 9.7% compared with planting distance of 25×10 cm. The plants that received N in three equal splits had higher grain yield (6993.1 kg.ha⁻¹). The Si-treated plants showed both higher yield parameters and greater physiological characteristics when compared with control plants. We observed an increase in grain yield (10%), N concentration (7%) and uptake (14.3%) in grain, protein content (6.8%), and nitrogen use efficiency (7.1%) by supplying Si fertilizer. Overall, our results revealed that foliar application of Si could be an effective technique for increasing rice grain yield and improving rice nutritional quality.

Keywords: rice yield, nutritional quality, silicon foliar application, planting space, plant nutrition.

INTRODUCTION

Rice (*Oryza sativa* L.) is one of the most important staple food crops in the world and is consumed by more than 50% of the world population (Zhou et al., 2018). Nitrogen (N) is one of the most important nutrients for the rice production and its deficiency or excess reduces the grain yield (Zhou et al., 2019). Proper management of N fertilizer usage in rice fields increases N use efficiency and improves grain yield in paddy fields and thus enhances the farmers' incomes in the regions (Faraji et al., 2012). N fertilizer should be applied at the right time to achieve the optimum grain yield (Thu et al., 2014). Therefore, N fertilization and optimal technique of N application are important for higher grain production (Sathiya and Ramesh, 2009). Proper application of N fertilizer is one of the most important

agronomic techniques that affect the yield and quality of rice (Lampayan et al., 2010). The increase in rice yield depends greatly on the N regulation at different growth stages (Pan et al., 2012). Optimal management of N application at different growth phases by regulating the source-sink relationship increases rice grain yield (Sui et al., 2013). Jafari Kelarijani et al. (2021) observed that efficient management of N fertilizer in paddy fields can be help to increase rice production potential. Kamruzzaman et al. (2013) suggested that N application in three equal splits at 15, 30 and 45 days after transplanting (DAT) would be essential for the production of high grain yield. In another study, Djaman et al. (2018) revealed that N application in four splits could be recommended to improve rice production and food security. Anil et al. (2018) found that the application of N (180 kg ha⁻¹) in four equal splits (basal,

20, 40 and 60 DAT) increased significantly the number of tillers m^{-2} and rice grain yield. Dahipahle and Singh (2018) reported that when N (150 kg ha^{-1}) was supply to rice in three splits as 33.3% at 15-20 days after sowing (DAS) + 33.3% at active tillering stage + 33.3% at panicle initiation stage enhanced significantly the crop growth and yield.

Among the agronomic techniques used to help rice plants to improve the grain yield and quality, is the foliar application of an abundant macro-element such as silicon (Si). Si is the second most abundant element in the Earth's crust after oxygen (Liao et al., 2020). Si is one of the beneficial elements for rice plants which can be increase the plant growth and productivity (Kheyri et al., 2019a). Si plays a vital role for rice plants by enhancing size and strength of sink (Lavinsky et al., 2016). It is documented that addition of Si to plant enhances the accumulation of nutrients such as Si, N, P and K in rice tissue (Cuong et al., 2017). The higher rice grain yield by Si foliar application could be attributed to greater nutrient accumulation in plant tissue (Kheyri et al., 2019b). Zia et al. (2017) found that the application of Si significantly increased the Si concentration in rice plants. Agostinho et al. (2017) demonstrated that the Si application can be an effective way for improving yield and increasing Si accumulation in rice. Patel et al. (2017) reported that Si nutrition helped to enhancing macronutrients and micronutrients content in rice grain and straw.

Choosing the right planting space (PS) is one of the vital factors in increasing the plant population (Tian et al., 2017). Reducing PS leads to an increase in plant density and ultimately affects the yield and quality of rice grains (Hu et al., 2020). Appropriate increase of plant population enhances the grain yield by balancing the yield components factors (Yang et al., 2019). Reduction of hill density leads to poor grain filling and thus reduces rice grain yield (Ao et al., 2019). Alipour Abookheili and Mobasser (2021) reported that the number of panicle m^{-2} and grain

yield in $30 \times 10 \text{ cm}^2$ PS were higher than $20 \times 20 \text{ cm}^2$ and $25 \times 25 \text{ cm}^2$ PS. In their 2-yr study, Sandhu et al. (2015) indicated that decreasing the planting distance increased grain yield. In a previous study on rice Kheyri et al. (2016) suggested the $20 \times 20 \text{ cm}$ PS as the suitable distance for Tarom Amrolahi variety. Therefore, determining the proper PS is necessary to improve grain yield and increase nitrogen use efficiency (NUE) (Zheng et al., 2020).

The aim of the present study was to investigate the effects of Si foliar application and N application time on yield components, grain yield, nutrient accumulation, protein content and N use efficiency in rice at different planting spaces.

MATERIAL AND METHODS

Experimental site

This experiment was performed at the farmer's field in Sari, Mazandaran Province, Iran ($36^{\circ}56'N$, $53^{\circ}01'E$; 15 m above sea level) during the two crop years of 2018 and 2019. The meteorological data of the experimental site during growth and development of rice for both crop years were provided from the nearest synoptic meteorological station to the paddy field and presented in Table 1. The soil of experimental site was classified as a Loam-textured. Soil sampling was performed at the experimental site at depths of 0-30 cm before the initiating experiment. Physical and chemical analyses of the soil included analysis of soil texture by hydrometric method (Bouyoucos, 1962), soil EC and pH by preparation of soil suspension (soil/water 1:5), organic matter using the volumetric method (Walkley and Black, 1934), total N by Kjeldahl method (Bremner and Mulvaney, 1982), extractable P by extraction sodium bicarbonate (Olsen et al., 1954), and extractable K by flame photometer techniques (Toth and Prince, 1949). Table 2 shows the soil physical and chemical properties of the experimental site.

SARVENAZ YAHYAZADEH ET AL.: IMPACTS OF SILICON FOLIAR SPRAYING
AND NITROGEN APPLICATION TECHNIQUES ON QUANTITATIVE AND QUALITATIVE PARAMETERS OF
RICE AT DIFFERENT PLANTING SPACES

Table 1. Meteorological parameters of the experimental site consisted of minimum (Min), maximum (Max) and average (Avg) temperature, total precipitation (TP) and total sunny hours monthly (TSHM) for both years (2018 and 2019)

| Months | Average temperature (°C) | | | | | | TP (mm) | | TSHM (h) | |
|--------|--------------------------|------|------|------|------|------|---------|------|----------|-------|
| | Min | | Max | | Avg | | 2018 | 2019 | 2018 | 2019 |
| | 2018 | 2019 | 2018 | 2019 | 2018 | 2019 | | | | |
| Apr | 10.3 | 10.6 | 21.3 | 18.8 | 15.8 | 14.7 | 38.3 | 78.2 | 148.7 | 131.5 |
| May | 15.1 | 15.3 | 25.7 | 25.8 | 20.4 | 20.6 | 18.6 | 51.1 | 194.5 | 200.6 |
| Jun | 20.0 | 21.4 | 28.3 | 31.7 | 24.1 | 26.5 | 12 | 1 | 171.2 | 272.4 |
| Jul | 24.5 | 24.0 | 34.5 | 32.0 | 29.5 | 28.0 | 7.4 | 69.9 | 255.4 | 202.4 |
| Aug | 25.2 | 23.3 | 32.9 | 32.3 | 29.0 | 27.8 | 72.6 | 22.3 | 142.8 | 184.2 |
| Sep | 24.5 | 20.6 | 31.7 | 29.4 | 28.1 | 25.0 | 11.9 | 45 | 182.8 | 160.3 |

Table 2. Physical and chemical characteristics of the experimental soil prior to experiment

| Parameters | Unit | Concentration | |
|----------------|---------------------|---------------|------|
| | | 2018 | 2019 |
| Sand | % | 39 | 42 |
| Silt | % | 36 | 32 |
| Clay | % | 25 | 26 |
| Textural class | | Loam | Loam |
| pH | | 7.04 | 7.54 |
| EC | ds m ⁻¹ | 0.86 | 1.41 |
| Organic matter | % | 2.82 | 2.70 |
| Total N | % | 0.18 | 0.16 |
| Available P | mg kg ⁻¹ | 7.9 | 8.1 |
| Available K | mg kg ⁻¹ | 165 | 181 |

Experimental design and treatments

The experiment was carried out as a split-split-plot in a randomized complete block design with three replications. The experimental treatments included the main plot assigned to planting space (25×10 cm, 20×10 cm), the sub-plot assigned to N application techniques (N application as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading, N application as 40% at basal + 40% at panicle initiation + 20% at full heading), and the sub-sub-plot assigned to foliar application of Si (control or non-use of Si, Si foliar application).

Rice cultivation and management

A local rice (*Oryza sativa* L.) cultivar Tarom Hashemi was used as plant material in the present study. The rice seeds were pregerminated by soaking in water for 24 h and incubated for 48 h in the dark. Germinated seeds were sown in nursery beds. The nursery land was prepared one week before sowing. Land preparation was

performed by puddling, harrowing and leveling the soil in the field. The plots size was 10 m² (2×5 m). The plots boundaries were enclosed with a plastic cover up to a depth of 30 cm to prevent the outflow of water and fertilizers between plots. All the experimental plots received phosphorus and potassium as basal at the rate of 130 kg P₂O₅ ha⁻¹ as triple superphosphate and 150 kg K₂O ha⁻¹ as potassium sulfate, respectively. N (urea 46%) was applied at the rate of 150 kg ha⁻¹ based on the treatments defined in the experimental plots. Si fertilizer from the source of potassium silicate (produced by AgriTecnico Company) was sprayed on plants at a rate of 5 per thousand. To control weeds, Butachlor herbicide was applied at a concentration of 3.5 L ha⁻¹ one week after transplanting, and the manual weeding was performed in week 2 and 4 after transplanting. In order to control *Chilo suppressalis*, diazinon insecticide (10% Granule) was applied at a rate of 15 kg ha⁻¹ at the maximum tillering and heading stages.

Sampling and measurement

At physiological maturity, plant samples were randomly selected from each plot after removing the border rows. The panicle length was determined by measuring 20 panicles in each plot. The number of total tillers hill⁻¹ was determined from 15 hills plot⁻¹. The number of panicles m⁻² was obtained by harvesting and counting all panicles from a 2 m² area in each plot. The Percentage of filled spikelets panicle⁻¹ was obtained from the ratio of the number of filled spikelets panicle⁻¹ to the total number of spikelets panicle⁻¹. The 1000-grain weight was recorded by counting and weighing 10 samples of 100 seeds. The grain yield were measured by manually harvesting an area of 4 m² (2×2 m) in the middle of each plot and based on 14% moisture content. The determination of accumulation of Si and N in rice grains was performed according to the methods described by Fallah et al. (2004) and Emami (1996), respectively. The N uptake in grain of rice was calculated by multiplying the dry matter accumulation in grain yield of rice with their respective concentrations expressed in kg ha⁻¹. The protein content in grain was calculated by total nitrogen multiplied by 5.95 (Lopez et al., 2010). The nitrogen use efficiency (NUE) was derived using the following formulae:

$$\text{NUE (kg.kg}^{-1}\text{)} = \frac{\text{grain yield (kg.ha}^{-1}\text{)}}{\text{nitrogen uptaked by the plant (kg.ha}^{-1}\text{)}}$$

Statistical analysis

Statistical analysis of data was performed using SAS software. Combined analysis of variance was conducted as split-split-plot in a randomized complete block design with three replications. Mean values were compared using least significant difference (LSD) test at 5% probability level.

RESULTS AND DISCUSSION

Growth parameters and grain yield

The results of combined analysis of variance showed that the main effect of year was significant on all the yield components. The grain yield was not affected by the year. The number of total tillers hill⁻¹, number of panicles⁻², and grain yield were significantly affected by the effect of planting space. The panicle length and grain yield were affected by the main effect of nitrogen application techniques. The simple effect of silicon was significant only on grain yield. Also, the 1000-grain weight was significantly (P≤0.05) affected by the three-way interaction between planting space × nitrogen application techniques × silicon foliar application (Table 3).

Table 3. Combined analysis of variance for PS, NAT and Si as well as their interactions on yield components and grain yield of rice

| Source of variation | df | Panicle length | No. of total tillers per plants | No. of panicle m ⁻² | Percent of filled spikelet panicle ⁻¹ | 1000-grain weight | Grain yield |
|---------------------------------------|----|---------------------|---------------------------------|--------------------------------|--|--------------------|--------------------------|
| Year (Y) | 1 | 10.82** | 210.50** | 632272.52** | 131.27* | 14.08** | 1208088.02 ^{ns} |
| Replication (Y) | 4 | 6.29 | 6.00 | 13820.83 | 16.13 | 1.16 | 1734897.39 |
| Planting space (PS) | 1 | 0.80 ^{ns} | 78.79** | 27888.52* | 18.32 ^{ns} | 0.75 ^{ns} | 5559004.68* |
| Y×PS | 1 | 3.21 ^{ns} | 5.89 ^{ns} | 20708.52 ^{ns} | 0.41 ^{ns} | 0.75 ^{ns} | 1471750.52 ^{ns} |
| Error | 4 | 2.63 | 2.86 | 4276.33 | 40.81 | 4.00 | 1671977.60 |
| Nitrogen application techniques (NAT) | 1 | 12.31** | 2.91 ^{ns} | 623.52 ^{ns} | 20.38 ^{ns} | 0.75 ^{ns} | 3799688.02* |
| Y×NAT | 1 | 13.83** | 4.38 ^{ns} | 999.18 ^{ns} | 13.04 ^{ns} | 0.08 ^{ns} | 2937825.52 ^{ns} |
| PS×NAT | 1 | 0.53 ^{ns} | 1.87 ^{ns} | 10296.02 ^{ns} | 5.86 ^{ns} | 10.08** | 174604.68 ^{ns} |
| Y×PS×NAT | 1 | 0.03 ^{ns} | 20.64* | 667.52 ^{ns} | 43.01* | 0.75 ^{ns} | 306400.52 ^{ns} |
| Error | 8 | 0.94 | 4.08 | 4217.87 | 17.04 | 1.91 | 1151534.37 |
| Silicon (Si) | 1 | 0.11 ^{ns} | 0.33 ^{ns} | 325.52 ^{ns} | 8.45 ^{ns} | 0.75 ^{ns} | 5751213.02* |
| Y×Si | 1 | 0.05 ^{ns} | 0.75 ^{ns} | 4780.02 ^{ns} | 2.86 ^{ns} | 0.08 ^{ns} | 519792.18 ^{ns} |
| PS×Si | 1 | 1.05 ^{ns} | 0.002 ^{ns} | 20542.68 ^{ns} | 46.49 ^{ns} | 0.75 ^{ns} | 400588.02 ^{ns} |
| Y×PS×Si | 1 | 0.29 ^{ns} | 2.53 ^{ns} | 16762.68 ^{ns} | 6.64 ^{ns} | 0.75 ^{ns} | 1838875.52 ^{ns} |
| NAT×Si | 1 | 4.56 ^{ns} | 4.28 ^{ns} | 13.02 ^{ns} | 3.95 ^{ns} | 2.08 ^{ns} | 17442.18 ^{ns} |
| Y×NAT×Si | 1 | 0.33 ^{ns} | 0.39 ^{ns} | 7326.02 ^{ns} | 20.38 ^{ns} | 0.08 ^{ns} | 129688.02 ^{ns} |
| PS×NAT×Si | 1 | 0.74 ^{ns} | 1.12 ^{ns} | 1598.52 ^{ns} | 2.52 ^{ns} | 4.08* | 88.02 ^{ns} |
| Y×PS×NAT×Si | 1 | 0.002 ^{ns} | 0.08 ^{ns} | 5963.02 ^{ns} | 27.63 ^{ns} | 0.08 ^{ns} | 1807692.18 ^{ns} |
| Error | 16 | 1.15 | 4.50 | 4662.81 | 29.21 | 0.58 | 817706.77 |
| CV (%) | - | 4.56 | 16.39 | 13.83 | 6.15 | 2.86 | 13.47 |

^{ns}, *, and ** are non-significant and significant at the 5 and 1% probability levels, respectively.

SARVENAZ YAHYAZADEH ET AL.: IMPACTS OF SILICON FOLIAR SPRAYING
AND NITROGEN APPLICATION TECHNIQUES ON QUANTITATIVE AND QUALITATIVE PARAMETERS OF
RICE AT DIFFERENT PLANTING SPACES

As shown in Table 4, the panicle length in the first year of the experiment was 4% higher than the second year. There was no significant difference in panicle length between two planting spaces, as well as between the foliar application and non-application of Si. Our results indicated that the panicle length increased by 4.2% when rice plants received the N fertilizer as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading compared with N application as 40% at basal

+ 40% at panicle initiation + 20% at full heading. These findings are in agreement with Kamruzzaman et al. (2013) and Alipour Abookheili et al. (2020), who reported that the N application in three equal splits significantly improved the rice panicle length. In this study, the panicle length was not affected by Si treatment. Similar to our results, the foliar application of Si had no significant effect on panicle length of rice (Sheykhzadeh et al., 2022).

Table 4. Mean comparison of main effects of Y, PS, NAT and Si on yield components and grain yield of rice

| Experimental treatments | Panicle length (cm) | No. of total tillers per plants | No. of panicle m ⁻² | Percent of filled spikelets panicle ⁻¹ | 1000-grain weight (g) | Grain yield (kg.ha ⁻¹) |
|---------------------------------|---------------------|---------------------------------|--------------------------------|---|-----------------------|------------------------------------|
| Year | | | | | | |
| First | 24.05a | 10.8b | 378.7b | 86.2a | 26.08b | 6553.1a |
| Second | 23.10b | 15.03a | 608.2a | 89.5a | 27.16a | 6870.4a |
| Planting space | | | | | | |
| 25×10 cm | 23.7a | 14.2a | 469.3b | 87.2a | 26.7a | 6371.5b |
| 20×10 cm | 23.4a | 11.6b | 517.5a | 88.4a | 26.5a | 7052.1a |
| Nitrogen application techniques | | | | | | |
| NAT1 | 24.08a | 12.6a | 489.8a | 88.5a | 26.7a | 6993.1a |
| NAT2 | 23.07b | 13.1a | 497.0a | 87.2a | 26.5a | 6430.4b |
| Silicon application | | | | | | |
| Control | 23.6a | 12.8a | 490.8a | 87.4a | 26.7a | 6365.6b |
| Silicon | 23.5a | 13.02a | 496.1a | 88.3a | 26.5a | 7057.9a |

Means in columns followed by the same letter(s) are not significantly different at $P \leq 0.05$;

NAT1: nitrogen application technique as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading;

NAT2: nitrogen application technique as 40% at basal + 40% at panicle initiation + 20% at full heading.

Our findings showed that the number of total tillers per plants in the second year was 28% higher than in the first year of the experiment. The higher growth parameters in the second year of this study could be attributed to improvement of environmental conditions, especially temperature and sunny hours (Table 1) during rice growing season compared with first year. It has been reported that there is a direct relationship between the yield components of rice and environmental factors such as light and temperature (Deng et al., 2015).

The results indicated that the increase in planting space (25×10 cm) led to an increase in number of total tillers by 18.3% compared with planting space of 20×10 cm (Table 4). The reduction in planting space followed by an increase in planting density reduced the number of tillers due to more competition

between rice seedlings for nutrient absorption (Alipour Abookheili and Mobasser, 2021). Our findings are further strengthened by the findings of Koireng et al. (2019), who reported that the number of tillers m⁻² significantly increased by enhancing planting spaces and reducing planting density. These findings are in agreement with Halder et al. (2018) and Kheyri et al. (2016). There was no significant difference between application and non-application of Si in terms of total tillers number. However, the Si foliar application had better positive impacts on number of total tillers than control plants. In similar results, Kheyri et al. (2019a) reported that the plants treated with both nano-Si and calcium silicate produced significantly higher total tillers when compared to control plants.

In the present study, the number of panicle m⁻² was higher in the second year than first

year by 37.7%, which was due to the increase in the total tillers number in the second year compared with first year. The Higher panicle m^{-2} results in greater grain yield and vice versa. We observed that the number of panicle m^{-2} significantly increased by decreasing planting space. The plants grown at 20×10 cm planting spacing showed higher panicle number m^{-2} (9.3%) when compared to plants grown at 25×10 cm spacing (Table 4). Our results are confirmed by Chen et al. (2019), who found that the number of panicle m^{-2} significantly increased by reduces in planting spaces and enhances in planting density. The findings in our study that the number of panicle m^{-2} exhibited a significant increase at 20×10 cm planting spacing was consistent with Kheyri et al. (2016) who reported that the number of panicle m^{-2} in rice significantly increased in response to decreasing planting spaces due to enhancing number of stems m^{-2} . Previous studies revealed that the panicle number m^{-2} plays a vital role in increasing rice grain yield (Kheyri et al., 2016; Chen et al., 2020). The Si treated plants had slight higher panicle number m^{-2} (496.1 panicles) over control plants (490.8 panicles), although there was no significant difference between application and non-use of Si. In another study, Sheykhzadeh et al. (2022) found that foliar application of Si at critical growth stages of rice helps to increase the total number of spikelets panicle⁻² and improve the rice grain yield.

Our results showed that there was no significant difference between various experimental treatments in terms of the percent of filled spikelets panicle⁻¹. However, the plants grown at 20×10 cm

planting spacing, N application in three equal splits (NAT1) and Si foliar application had positive effects on percent of filled spikelets panicle⁻¹ (Table 4). Alipour Abookheili et al. (2020) were observed the higher number of filled spikelets when N fertilizer applied in three equal splits. Kheyri et al. (2019a, b) demonstrated that Si-treated plants had higher number of filled grains panicle⁻¹ when compared with plants did not receive Si fertilizer. Lavinsky et al. (2016) reported that the Si application (2 mM) at reproductive growth phase of rice enhanced the number of filled grains panicle⁻¹.

The results presented in Table 5 indicated that the 1000-grain weight had higher value (27.66 g) at planting distance of 25×10 cm by Si foliar spraying and N application as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading, whereas the plants grown at planting spaces of 20×10 cm showed higher thousand-grain weight (TGW) (27.33 g) when Si foliar sprayed and N fertilizer applied as 40% at basal + 40% at panicle initiation + 20% at full heading. This result reveals that at planting distance of 20×10 cm, more N fertilizer application at the vegetative growth stage along with Si application can be help to ameliorate TGW weight. However, the greatest TGW in both planting spaces was observed in Si-treated plants. Kheyri et al. (2019a, b) also observed that Si application via either nano-Si or calcium silicate improved growth and yield of rice plants compared with control plants. In a 2-yr study, Dahipahle and Singh (2018) found that the N application in three equal splits at the vegetative and reproductive growth phases resulted in highest TGW.

Table 5. Mean comparison of interactions between PS, NAT and Si on 1000-grain weight of rice

| Planting space | 25×10 cm | | 20×10 cm | |
|---------------------|----------|----------|-----------|----------|
| | NAT1 | NAT2 | NAT1 | NAT2 |
| Treatments | | | | |
| Control | 27.00abc | 26.00cd | 26.66abcd | 26.33bcd |
| Silicon application | 27.66a | 26.33bcd | 25.66d | 27.33ab |

Means in columns followed by the same letter(s) are not significantly different at $P \leq 0.05$;

NAT1: nitrogen application technique as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading;

NAT2: nitrogen application technique as 40% at basal + 40% at panicle initiation + 20% at full heading.

SARVENAZ YAHYAZADEH ET AL.: IMPACTS OF SILICON FOLIAR SPRAYING
AND NITROGEN APPLICATION TECHNIQUES ON QUANTITATIVE AND QUALITATIVE PARAMETERS OF
RICE AT DIFFERENT PLANTING SPACES

The assay of grain yield revealed that the plants grown at planting distance of 20×10 cm had highest grain yield (7052.1 kg.ha⁻¹), whereas the increase in planting space (25×10 cm) resulted in the yield reduction of 9.7% (Table 4). The higher yield at planting space of 20×10 cm could be attributed to greater panicle number m⁻². Alipour Abookheili et al. (2020) indicated that reducing planting space from 25×25 cm to 30×10 cm enhanced rice grain yield by 20.3% by increasing number of panicles m⁻². Similar findings were confirmed by Zhou et al. (2019), Zheng et al. (2020) and Chen et al. (2019), who reported that higher grain yield was observed in lower planting spaces followed by higher planting density.

In this research, the application of N fertilizer in three equal splits (NAT1) significantly increased the grain yield by 8% when compared with NAT2 treatment (Table 4). Optimal splitting of N fertilizer at different growth stages improves rice grain yield by enhancing N uptake by the plant (Esmailzadeh Moridani et al., 2011). Faraji et al. (2011) documented the increase in grain yield of rice by N splitting application in appropriate doses.

We also observed that the plants untreated with Si showed the lower yield (6365.6 kg.ha⁻¹), whereas the Si-treated plants had 10% higher

grain yield than control plants (Table 4). Foliar application of Si increases the rice grain yield by facilitating the nutrients uptake by the plant (Kheyri et al., 2018). Cuong et al. (2017) reported that the Si-treated rice plants showed higher grain yield due to increasing yield attributes and ameliorating nutrients accumulation. Kheyri et al. (2019a) found that the application of calcium silicate and nano-Si increased rice grain yield compared with control plants by 6.9% and 9.6%, respectively, which is consistent with the results of the present study.

Physiological parameters

The results of combined analysis of variance revealed that the all physiological parameters were significantly affected by the simple effects of year. The main effects of planting spaces and nitrogen application techniques were not significant on physiological characteristics of rice. Among physiological traits, only nitrogen uptake in grain was significantly ($P < 0.05$) affected by silicon foliar application. The physiological parameters were not affected by the three-way interaction between planting space × nitrogen application techniques × silicon foliar application (Table 6).

Table 6. Combined analysis of variance for PS, NAT and Si as well as their interactions on Si and N accumulation in grain, protein content in grain and nitrogen use efficiency of rice

| Source of variation | df | Si concentration in grain | N concentration in grain | N uptake in grain | Protein content in grain | N Use Efficiency |
|---------------------------------------|----|---------------------------|--------------------------|-----------------------|--------------------------|----------------------|
| Year (Y) | 1 | 0.229** | 0.569** | 1708.10* | 20.30** | 868.45** |
| Replication (Y) | 4 | 0.012 | 0.096 | 1073.81 | 3.42 | 8.48 |
| Planting space (PS) | 1 | 0.016 ^{ns} | 0.048 ^{ns} | 276.16 ^{ns} | 1.74 ^{ns} | 253.12 ^{ns} |
| Y×PS | 1 | 0.001 ^{ns} | 0.011 ^{ns} | 53.19 ^{ns} | 0.42 ^{ns} | 21.10 ^{ns} |
| Error | 4 | 0.039 | 0.034 | 515.85 | 1.22 | 32.14 |
| Nitrogen application techniques (NAT) | 1 | 0.010 ^{ns} | 0.084 ^{ns} | 3.66 ^{ns} | 2.99 ^{ns} | 242.82 ^{ns} |
| Y×NAT | 1 | 0.006 ^{ns} | 0.065 ^{ns} | 1197.15 ^{ns} | 2.29 ^{ns} | 19.01 ^{ns} |
| PS×NAT | 1 | 0.003 ^{ns} | 0.035 ^{ns} | 372.63 ^{ns} | 1.25 ^{ns} | 54.53 ^{ns} |
| Y×PS×NAT | 1 | 0.007 ^{ns} | 0.036 ^{ns} | 58.60 ^{ns} | 1.29 ^{ns} | 143.77 ^{ns} |
| Error | 8 | 0.018 | 0.032 | 262.37 | 1.14 | 40.28 |
| Silicon (Si) | 1 | 0.002 ^{ns} | 0.089 ^{ns} | 1951.80* | 3.15 ^{ns} | 60.23 ^{ns} |
| Y×Si | 1 | 0.011 ^{ns} | 0.312* | 2814.65** | 11.06* | 60.62 ^{ns} |
| PS×Si | 1 | 0.006 ^{ns} | 0.002 ^{ns} | 89.21 ^{ns} | 0.08 ^{ns} | 2.18 ^{ns} |
| Y×PS×Si | 1 | 0.001 ^{ns} | 0.222 ^{ns} | 275.61 ^{ns} | 7.93 ^{ns} | 111.59 ^{ns} |
| NAT×Si | 1 | 0.038 ^{ns} | 0.062 ^{ns} | 373.49 ^{ns} | 2.20 ^{ns} | 8.48 ^{ns} |
| Y×NAT×Si | 1 | 0.30 ^{ns} | 0.000 ^{ns} | 55.61 ^{ns} | 0.001 ^{ns} | 6.50 ^{ns} |
| PS×NAT×Si | 1 | 0.004 ^{ns} | 0.000 ^{ns} | 1.55 ^{ns} | 0.01 ^{ns} | 10.76 ^{ns} |
| Y×PS×NAT×Si | 1 | 0.000 ^{ns} | 0.007 ^{ns} | 117.27 ^{ns} | 0.25 ^{ns} | 64.15 ^{ns} |
| Error | 16 | 0.024 | 0.065 | 328.45 | 2.32 | 68.24 |
| CV (%) | - | 15.80 | 20.67 | 21.91 | 20.68 | 27.35 |

^{ns}, *, and ** are non-significant and significant at the 5 and 1% probability levels, respectively.

In this study, the Si grain concentration in the second year (1.05%) was higher than first year (0.91%). Our findings showed that the different experimental treatments did not have a significant effect on Si concentration in rice grains. However, the plants grown at planting space of 20×10 cm, N application as 40% at basal + 40% at panicle initiation + 20% at full heading and foliar application of Si on plants produced higher Si concentration in grain (Table 7). Razavipour et al. (2018) observed the positive impacts of N application

at different growth phases of rice plants on nutrients uptake in grain. The correct technique of using N fertilizer during the rice growing season can be helps to enhancing plant growth, improving photosynthesis, and increasing N concentration in grain (Moslehi et al., 2016). Our results are in agreement with Kheyri et al. (2019a) who documented that addition of Si to rice plants enhanced the grain Si accumulation. Cuong et al. (2017) also observed the higher Si content in rice grains by application of Si fertilizer.

Table 7. Mean comparison of main effects of Y, PS, NAT and Si on N and Si accumulation in grain, protein content in grain and nitrogen use efficiency of rice

| Experimental treatments | Si concentration in grain (%) | N concentration in grain (%) | N uptake in grain (kg.ha ⁻¹) | Protein content in grain (%) | N Use Efficiency (kg.kg ⁻¹) |
|---------------------------------|-------------------------------|------------------------------|--|------------------------------|---|
| Year | | | | | |
| First | 0.91b | 1.34a | 88.67a | 8.01a | 25.95b |
| Second | 1.05a | 1.12b | 76.74b | 6.71b | 34.45a |
| Planting space | | | | | |
| 25×10 cm | 0.96a | 1.27a | 80.30a | 7.55a | 27.90a |
| 20×10 cm | 1.00a | 1.20a | 85.10a | 7.17a | 32.50a |
| Nitrogen application techniques | | | | | |
| NAT1 | 0.96a | 1.19a | 82.98a | 7.11a | 32.45a |
| NAT2 | 0.99a | 1.28a | 82.43a | 7.61a | 27.95a |
| Silicon application | | | | | |
| Control | 0.97a | 1.19a | 76.33b | 7.10a | 29.08a |
| Silicon | 0.98a | 1.28a | 89.08a | 7.62a | 31.32a |

Means in columns followed by the same letter(s) are not significantly different at $P \leq 0.05$;

NAT1: nitrogen application technique as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading;

NAT2: nitrogen application technique as 40% at basal + 40% at panicle initiation + 20% at full heading.

The results presented in Table 7 showed that the N concentration in grain in the first year (1.34%) was significantly higher than the second year (1.12%) of this study. We found that the grain N uptake in plants grown in the first year of the experiment had a similar trend to grain N concentration. The planting space of 20×10 cm resulted in 6% higher N uptake than planting space of 25×10 cm. In this research, the grain N uptake was increased in response to Si fertilization. The Si foliar application enhanced N uptake in rice grains by 14.3% compared with control plants. The augmentation in growth, yield and quality of rice could be attributed to the ameliorating uptake of nutrients such as Si, Zn and N due to the application of Si fertilizer (Kheyri et al., 2019a). Gou et al. (2020) documented the positive impacts of Si on N nutritional processes such as uptake

such as uptake, assimilation, and remobilization. Si is able to improve N uptake in rice plants in the situation of non-optimal N supply (Deus et al., 2020). Liang et al. (2021) found that the proper application of Si fertilizer helped to improving N availability for rice plants in paddy fields. Our findings are confirmed by Patel et al. (2017) and Cuong et al. (2017) who demonstrated that Si application increased the N accumulation in rice plant tissue.

The protein content was only affected by the year. Likewise, it was significantly higher in the first year (8.01%) than in the second year (6.71%) (Table 7). The higher protein content in the first year was due to the greater N concentration and uptake in this year compared with second year. N fertilizer affects rice grain quality by increasing protein accumulation (Hao et al., 2007). In

this study, the use of Si had slight better impacts on grain protein content compared to non-application of Si. Yazdpour et al. (2014) suggested that foliar application of Si on rice plants significantly increased N concentration and protein content, which is consistent with the results of the present study. Liu et al. (2017) indicated that addition of Si to rice significantly increased the protein content by 6.19% and 5.52% in milled and brown rice, respectively. Our results are further strengthened by the findings of Zeng et al. (2011), Cuong et al. (2017) and Kheyri et al. (2019a), who observed the enhancing protein content and improving grain quality in rice by supplying Si fertilizer.

As shown in Table 7, the nitrogen use efficiency (NUE) in the second year was 24.7% higher than in the first year. The results indicated that the NUE was higher at planting space of 20×10 cm, N application as 33.3% at basal + 33.3% at panicle initiation + 33.3% at full heading, and foliar application of Si by 14.1%, 13.9% and 7.1%, compared with planting space of 25×10 cm, N application as 40% at basal + 40% at panicle initiation + 20% at full heading, and non-use of Si fertilizer, respectively. It has been reported that the N physiological efficiency in rice reduced by the increase of N application rate (Yesuf and Balcha, 2014). Dehpouri et al. (2022) found that the increase in N consumption resulted in the reduction of N efficiency. On the other hand, Asadi et al. (2012) mentioned that the NUE was not affected by N application in different rice growth stages. Si application increases N uptake and NUE by improving source capacity and sink strength (Mohanty et al., 2019). Sabaghnia et al. (2018) reported that the Si-treated plants showed higher fertilizer efficiency compared with control plants. Our results are in agreement with Liao et al. (2020) who documented that Si application could be an effective technique for increasing plant growth and improving N fertilizer efficiency. Increasing NUE at whole-plant level of wheat by using Si fertilizer has also been reported (Neu et al., 2017).

CONCLUSIONS

Our findings illustrated that the plants grown at planting space of 20×10 cm showed the higher panicles number m^{-2} , followed by higher grain yield, and greater N uptake in grain and higher NUE compared with 25×10 cm planting spacing. The NAT1 treatment improved the grain yield and NUE when compared with NAT2 treatment. The addition of Si fertilizer to rice plants ameliorated the yield components, grain yield, and physiological parameters of rice compared with control plants. The foliar application of Si was able to enhance the yield by 10%, improve the N uptake by 14.3%, and increase the NUE by 7.1% in rice plants compared with non-application of Si. Also, an increase in the concentrations of Si and N, and an improvement in the protein content were observed for rice receiving foliar Si. Overall, the results of the present study indicated that the foliar application of Si fertilizer could increase rice grain yield by enhancing yield components and improve rice grain quality by increasing nutrients accumulation and protein content in rice.

REFERENCES

- Agostinho, F.B., Tubana, B.S., Martins, M.S., Datnoff, L.E., 2017. *Effect of different silicon sources on yield and silicon uptake of rice grown under varying phosphorus rates*. *Plants*, 6: 35.
- Alipour Abookheili, F., Noormohammadi, G., Madani, H., Heidari Sharifabad, H., Mobasser, H., 2020. *Effect of nitrogen splitting and plant density on yield and grain yield components of two rice genotypes (Oryza sativa L.)*. *Iranian Journal of Field Crops Research*, 17(4): 631-645.
- Alipour Abookheili, F., and Mobasser, H.R., 2021. *Effect of planting density on growth characteristics and grain yield increase in successive cultivations of two rice cultivars*. *Agrosystems, Geosciences and Environment*, 4: e20213.
- Anil, K., Yakadri, M., Jayasree, G., 2018. *Influence of nitrogen levels and times of application on growth parameters of aerobic rice*. *International Journal of Current Microbiology and Applied Sciences*, 7 (5): 1525-1529.
- Ao, H., Xie, X., Huang, M., Zou, Y., 2019. *Decreasing hill density combined with increasing nitrogen rate*

- led to yield decline in hybrid rice under low-light conditions. *Scientific Reports*, 9: 15786.
- Asadi, S., Zavareh, M., Shokri Vahed, H., Shahin Rokhsar, P., 2012. *Effect of supplement foliar application of nitrogen and potassium on yield, grain quality and nitrogen utilization efficiency of hybrid rice c.v. Bahar-1*. *Journal of Crop Production*, 4(3): 175-190.
- Bouyoucos, G.J., 1962. *Hydrometer method improved for making particle size analysis of soils*. *Agronomy Journal*, 54: 464-465.
- Bremner, J.M., and Mulvaney, C.S., 1982. *Nitrogen-Total*. In: Page, A.L., Miller, R.H., Keeney D.R. (eds.), *Methods of soil analysis, Part 2, Chemical and microbiological properties*. *Agronomy Monographs*, ASA, SSSA, Madison, Wisconsin, USA: 595-624.
- Chen, S., Yin, M., Zheng, X.I., Liu, S., Chu, G., Xu, C., Wang, D., Zhang, X., 2019. *Effect of dense planting of hybrid rice on grain yield and solar radiation use in southeastern China*. *Agronomy Journal*, 111: 1229-1238.
- Chen, J., Zhang, R., Cao, F., Yin, X., Zou, Y., Huang, M., Abou-Elwafa, S.F., 2020. *Evaluation of late-season short- and long-duration rice cultivars for potential yield under mechanical transplanting conditions*. *Agronomy*, 10: 1307.
- Cuong, T.X., Ullah, H., Datta, A., Hanh, T.C., 2017. *Effects of silicon-based fertilizer on growth, yield and nutrient uptake of rice in tropical zone of Vietnam*. *Rice Science*, 24(5): 283-290.
- Dahipahle, A.V., and Singh, U.P., 2018. *Effect of crop establishment, nitrogen levels and time of nitrogen application on growth and yield attributing parameters of direct seeded rice (Oryza sativa L.)*. *International Journal of Chemical Studies*, 6(2): 2889-2893.
- Dehpouri, F., Barari Tari, D., Niknejad, Y., Fallah Amoli, H., Amiri, E., 2022. *Study of nitrogen fertilization management on corn yield and nitrogen use efficiency in the southern Caspian Sea region*. *Romanian Agricultural Research*, 39: 385-390.
- Deng, N., Ling, X., Sun, Y., Zhang, C., Fahad, S., Peng, S., Cui, K., Nie, L., Huang, J., 2015. *Influence of temperature and solar radiation on grain yield and quality in irrigated rice system*. *European Journal of Agronomy*, 64: 37-46.
- Deus, A.C.F., de Mello Prado, R., de Cássia Félix Alvarez, R., de Oliveira, R.L.L., Felisberto, G., 2020. *Role of silicon and salicylic acid in the mitigation of nitrogen deficiency stress in rice plants*. *Silicon*, 12: 997-1005.
- Djaman, K., Mel, V.C., Ametonou, F.Y., El-Namaky, R., Diallo, M.D., Koudahe, K., 2018. *Effect of nitrogen fertilizer dose and application timing on yield and nitrogen use efficiency of irrigated hybrid rice under semi-arid conditions*. *Journal of Agricultural Science and Food Research*, 9: 223.
- Emami, A., 1996. *Methods of plant analysis. Vol. 982*. Soil and Water Res. Institute, Tehran, Iran.
- Esmailzadeh Moridani, M., Eshraghi-Nejad, M., Galeshi, S., Ashouri, M., 2011. *The investigation of nitrogen fertilizer split application effect on quantity yield and grain quality of rice varieties (Hashemi and Bahar 1) in Guilan*. *Electronic Journal of Crop Production*, 4(2): 121-137.
- Fallah, A., Visperas, R.M., Alejar, A.A., 2004. *The interactive effect of silicon and nitrogen on growth and spikelet filling in rice (Oryza sativa L.)*. *Philippine Agricultural Scientist*, 87: 174-176.
- Faraji, F., Esfahani, M., Kavooosi, M., Nahvi, M., Rabiei, B., 2011. *Effect of nitrogen fertilizer application on grain yield and milling recovery of rice (Oryza sativa cv. Khazar)*. *Iranian Journal of Crop Science*, 13(1): 61-77.
- Faraji, F., Esfehiani, M., Kavooosi, M., Nahvi, M., Rabiyyi, B., 2012. *Effects of split application and levels of nitrogen fertilizer on growth indices and grain yield of rice (Oryza sativa Cv. Khazar)*. *Iranian Journal of Field Crop Science*, 43(2): 323-333.
- Gou, T., Yang, L., Hu, W., Chen, X., Zhu, Y., Guo, J., Gong, H., 2020. *Silicon improves the growth of cucumber under excess nitrate stress by enhancing nitrogen assimilation and chlorophyll synthesis*. *Plant Physiology and Biochemistry*, 152: 53-61.
- Halder, J., Rokon, G.M., Islam, M.A., Salahin, N., Alam, M.K., 2018. *Effect of planting density on yield and yield attributes of local aromatic rice varieties*. *Bangladesh Journal of Agricultural Research*, 43: 489-497.
- Hao, H.L., Wei, Y.Z., Yang, X.E., Feng, Y., Wu, C.Y., 2007. *Effects of different nitrogen fertilizer levels on Fe, Mn, Cu and Zn concentrations in Shoot and grain quality in rice (Oryza sativa)*. *Rice Science*, 14: 289-294.
- Hu, Q., Jiang, W.Q., Qiu, S., Xing, Z.P., Hu, Y.J., Guo, B.W., Liu, G.D., Gao, H., Zhang, H.C., Wei, H.Y., 2020. *Effect of wide-narrow row arrangement in mechanical pot-seedling transplanting and plant density on yield formation and grain quality of japonica rice*. *Journal of Integrative Agriculture*, 19(5): 1197-1214.
- Jafari Kelarijani, S.M., Barari Tari, D., Niknejad, Y., Fallah, H., Amiri, E., 2021. *Nitrogen affects on rice growth and nitrogen efficiency indices in different geographical regions in northern Iran*. *Romanian Agricultural Research*, 38: 79-92.
- Kamruzzaman, M.D., Abdul Kayum, M.D., Mainul Hasan, M.D., Mahmudul Hasan, M.D., Da Silva, J.A.T., 2013. *Effect of split application of nitrogen fertilizer on yield and yield attributes of transplanted aman rice (Oryza sativa L.)*. *Bangladesh Journal of Agricultural Research*, 38(4): 579-587.
- Kheyri, N., Mobasser, H.R., Masoodi, B., Yadollahi, P., 2016. *Effect of plant density and planting pattern on yield components and yield of rice (Oryza sativa L.) var. Tarom Amrollahi*. *Journal of Plant Ecophysiology*, 8: 26-34.

SARVENAZ YAHYAZADEH ET AL.: IMPACTS OF SILICON FOLIAR SPRAYING
AND NITROGEN APPLICATION TECHNIQUES ON QUANTITATIVE AND QUALITATIVE PARAMETERS OF
RICE AT DIFFERENT PLANTING SPACES

- Kheyri, N., Ajam Norouzi, H., Mobasser, H.R., Torabi, B., 2018. *Effect of different resources and methods of silicon and zinc application on agronomic traits, nutrient uptake and grain yield of rice (Oryza sativa L.)*. Applied Ecology and Environmental Research, 16: 5781-5798.
- Kheyri, N., Ajam Norouzi, H., Mobasser, H.R., Torabi, B., 2019a. *Effects of silicon and zinc nanoparticles on growth, yield, and biochemical characteristics of rice*. Agronomy Journal, 111: 3084-3090.
- Kheyri, N., Ajam Norouzi, H., Mobasser, H.R., Torabi, B., 2019b. *Comparison of NPs foliar application of silicon and zinc with soil application on agronomic and physiological traits of rice (Oryza sativa L.)*. Iranian Journal of Field Crops Research, 17: 503-515.
- Koireng, R.J., Devi, M.N.G., Devi, P.K.H., Gogoi, M., Rolling Anal, P.S., 2019. *Effect of variety and spacing on the productivity of direct seeded rice (Oryza sativa L.) under Manipur condition*. Indian Journal of Pure and Applied Biosciences, 7: 335-341.
- Lampayan, R.M., Bouman, B.A.M., Dios, J.L.D., Espirity, A.J., Soriano, J.B., Lactaen, A.T., Faronilo, J.E., Thant, K.M., 2010. *Yield of aerobic rice in rainfed lowlands of the Philippines as affected by nitrogen management and row spacing*. Field Crops Research, 116: 165-174.
- Lavinsky, A.O., Detmann, K.C., Reis, J.V., Ávila, R.T., Sanglard, M.L., Pereira, L.F., DaMatta, F.M., 2016. *Silicon improves rice grain yield and photosynthesis specifically when supplied during the reproductive growth stage*. Journal of Plant Physiology, 206: 125-132.
- Liang, Y., Liao, M., Fang, Z., Guo, J., Xie, X., Xu, C., 2021. *How silicon fertilizer improves nitrogen and phosphorus nutrient availability in paddy soil?* Journal of Zhejiang University - Science B (Biomedicine and Biotechnology), 22(7): 521-532.
- Liao, M., Fang, Z., Liang, Y., Huang, X., Yang, X., Chen, S., Xie, X., Xu, C., Guo, J., 2020. *Effects of supplying silicon nutrient on utilization rate of nitrogen and phosphorus nutrients by rice and its soil ecological mechanism in a hybrid rice double-cropping system*. Journal of Zhejiang University - Science B (Biomedicine and Biotechnology), 21: 474-484.
- Liu, Q., Zhou, X., Sun, Z., 2017. *Application of silicon fertilizer affects nutritional quality of rice*. Chilean Journal of Agricultural Research, 77(2): 163-170.
- Lopez, C.V.G., Garcia, M.D.C.C., Fernandez, F.G.A., Bustos, C.S., Chisti, Y., Sevilla, J.M.F., 2010. *Protein measurements of microalgal and cyanobacterial biomass*. Bioresource Technology, 101: 7587-7591.
- Mohanty, S., Nayak, A.K., Swain, C.K., Dhal, B., Kumar, A., Tripathi, R., Shahid, M., Lal, B., Gautam P., Dash, G.K., Swain, P., 2019. *Silicon enhances yield and nitrogen use efficiency of tropical low land rice*. Agronomy Journal, 112: 70-84.
- Moslehi, N., Niknejad, Y., Fallah Amoli, H., Kheyri, N., 2016. *Effect of integrated application of chemical, organic and biological fertilizers on some of the morphophysiological traits of rice (Oryza sativa L.) Tarom Hashemi cultivar*. Crop Physiology Journal, 8(30): 87-103.
- Neu, S., Schaller, J., Dudel, E.G., 2017. *Silicon availability modifies nutrient use efficiency and content, C:N:P stoichiometry, and productivity of winter wheat (Triticum aestivum L.)*. Scientific Reports, 7: 40829.
- Olsen, S.R., Cole, C.V., Watanabe, F.S., Dean, L.A., 1954. *Estimation of available phosphorus in soils by extraction with sodium bicarbonate*. Circ. 939, USDA, Washington DC.
- Pan, S.G., Huang, S.Q., Zhai, J., Wang, J.P., Cao, C.G., Cai, M.L., Zhan, M., Tang, X.R., 2012. *Effects of N management on yield and N uptake of rice in central China*. Journal of Integrative Agriculture, 11: 1993-2000.
- Patel, R.A., Patel, K.C., Malav, J.K., 2017. *Status of silicon in rice (Oryza sativa L.) and its correlation with other nutrients under Typic ustochrepts soil*. International Journal of Current Microbiology and Applied Sciences, 6(12): 2598-2611.
- Razavipour, T., Khaledian, M., Rezaei, M., 2018. *Effects of nitrogen levels and its splitting on rice yield and nutrient uptake in rice, Hashemi variety*. Human Environment, 45: 153-164.
- Sabaghnia, N., Javanmard, A., Janmohammadi, M., Nouraein, M., 2018. *The influence of nano-TiO₂ and nano-Silica particles effects on yield and morphological traits of sunflower*. Helia, 41: 213-225.
- Sandhu, S.S., Mahal, S.S., Kaur, A., 2015. *Physicochemical, cooking quality and productivity of rice as influenced by planting methods, planting density and nitrogen management*. International Journal of Food, Agriculture and Veterinary Sciences, 5(1): 33-40.
- Sathiya, K., and Ramesh, T., 2009. *Effect of split application of nitrogen on growth and yield of aerobic rice*. Asian Journal of Experimental Sciences, 23 (1): 303-306.
- Sheykhzadeh, M., Mobasser, H., Rahimi Petrodi, E., Rezvani, M., 2022. *Silicon and zinc improves grain yield and nutrient status in rice when supplied during the different growth stages*. Romanian Agricultural Research, 39: 371-383.
- Sui, B., Feng, X.M., Tian, G.L., Hu, X.Y., Shen, Q.R., Guo, S.W., 2013. *Optimizing nitrogen supply increases rice yield and nitrogen use efficiency by regulating yield formation factors*. Field Crop Research, 150: 99-107.
- Thu, T.T.P., Yamakawa, T., Moe, K., 2014. *Effect of nitrogen application timing on growth, grain yield and eating quality of the KD18 and TH3-3 rice varieties*. Journal of the Faculty of Agriculture, Kyushu University, 59 (1): 55-64.

- Tian, G., Gao, L., Kong, Y., Hu, X., Xie, K., Zhang, R., Ling, N., Shen, Q., Guo, S., 2017. *Improving rice population productivity by reducing nitrogen rate and increasing plant density*. PLoS One, 12(8): e0182310.
- Toth, S.J., and Prince, A.L., 1949. *Estimation of cation exchange capacity and exchangeable Ca, K and Na contents of soils by flame photometer techniques*. Soil Science, 67: 439-446.
- Walkley, A., Black, I.A., 1934. *Estimation of soil organic carbon by the chromic acid titration method*. Soil Science, 37: 29-38.
- Yang, D., Cai, T., Luo, Y., Wang, Z., 2019. *Optimizing plant density and nitrogen application to manipulate tiller growth and increase grain yield and nitrogen-use efficiency in winter wheat*. PeerJ, 7: e6484.
- Yazdpour, H., Noormohamadi, G., Madani, H., Heidari Sharif Abad, H., Mobasser, H.R., Oshri, M., 2014. *Role of nano-silicon and other silicon resources on straw and grain protein, phosphorus and silicon contents in Iranian rice cultivar (Oryza sativa cv. Tarom)*. International Journal of Biosciences, 5(12): 449-456.
- Yesuf, E., and Balcha, A., 2014. *Effect of nitrogen application on grain yield and nitrogen efficiency of rice (Oryza sativa L.)*. Asian Journal of Crop Science, 6(3): 273-280.
- Zeng, F.R., Zhao, F.S., Qiu, B.Y., Quyang, Y.N., Wu, F.B., Zhang, G.P., 2011. *Alleviation of chromium toxicity by silicon addition in rice plants*. Agricultural Sciences in China, 10: 1188-1196.
- Zheng, H., Chen, Y., Chen, Q., Li, B., Zhang, Y., Jia, W., Mo, W., Tang, Q., 2020. *High-density planting with lower nitrogen application increased early rice production in a double-season rice system*. Agronomy Journal, 112: 205-214.
- Zhou, C.C., Huang, Y.C., Jia, B.Y., Wang, Y., Wang, Y., Xu, Q., Li, R.F., Wang, S., Dou, F.G., 2018. *Effects of cultivar, nitrogen rate, and planting density on rice-grain quality*. Agronomy, 8: 1-13.
- Zhou, C., Huang, Y., Jia, B., Wang, S., Dou, F., Samonte, S.O.P.B., Chen, K., Wang, Y., 2019. *Optimization of nitrogen rate and planting density for improving the grain yield of different rice genotypes in northeast China*. Agronomy, 9: 555.
- Zia, Z., Bakhat, H.F., Saqib, Z.A., Shah, G.M., Fahad, S., Ashraf, M.R., Hammad, H.M., Naseem, W., Shahid, M., 2017. *Effect of water management and silicon on germination, growth, phosphorus and arsenic uptake in rice*. Ecotoxicology and Environmental Safety, 144: 11-18.